

Rehabilitation Protocol: Knee Arthroscopy Meniscectomy/Chondroplasty/Synovectomy

| Name: | | Date: |
|---------------------------------------|--|---------------------------------------|
| Diagnosis: | | Date of Surgery: |
| Phase I (Wee | ks 0-2) | |
| RangeThera | ntbearing: As tolerated with crutches (for balance of Motion – AROM as tolerated Goal: Immediate full range of motion peutic Exercises Quad and Hamstring sets Heel slides | nce) x 24-48 hours – progress to WBAT |
| 0 0 | Co-contractions Isometric adduction and abduction exercises Straight-leg raises Patellar mobilization | |
| Phase II (We | eks 3-5) | |
| Range Thera | ntbearing: As tolerated e of Motion – maintain full ROM – gentle passiv peutic Exercises Quadriceps and Hamstring strengthening Lunges Wall-sits Balance exercises – Core work | e stretching at end ranges |
| Phase III (Weeks 6-8) | | |
| RangeThera | e of Motion – Full/Painless ROM peutic Exercises Leg press Hamstring curls Squats Plyometric exercises Endurance work Return to athletic activity as tolerated | |
| Comments: | | |
| Frequency: 2-3 times per week | | Duration: 6-8 weeks |
| Signature: | | Date: |



