

## Rehabilitation Protocol: Pectoralis Major Repair

<ul> <li>hand exercises, patient may ride stationary bike with operative arm in the sling</li> <li>Phase II: 4-6 Weeks</li> <li>Sling Immobilizer: Worn at all times - Sleep with pillow under elbow to support the operative arm</li> <li>Range of Motion: AAROM in the supine position with wand - Goal: Forward Flexion to 90 degrees</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, shoulder shrugs/scapular retraction without resistance</li> <li>Phase III: 7-9 Weeks</li> <li>Sling Immobilizer: May be discontinued</li> <li>Range of Motion: AROM in the pain free range - No PROM, AAROM (pulleys, supine wand, wall clin)</li> <li>Goals: Forward Flexion to 120 degrees and Abduction to 90 degrees, ER to tolerance, IR and E ension (wand behind the back)</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, submaximal isometrics</li> <li>Phase IV: 10-12 Weeks</li> <li>Range of Motion: AROM and AAROM in the pain free range - No PROM, Goals: Full ROM</li> </ul>	Name:		Date:		
<ul> <li>Sling Immobilizer: Worn at all times - Sleep with pillow under elbow to support the operative arm         <ul> <li>Range of Motion: Supported pendulum exercises under guidance of PT</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, patient may ride stationary bike with operative arm in the sling</li> </ul> </li> <li>Phase II: 4 - 6 Weeks         <ul> <li>Sling Immobilizer: Worn at all times - Sleep with pillow under elbow to support the operative arm</li> <li>Range of Motion: AAROM in the supine position with wand - Goal: Forward Flexion to 90 degrees</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, shoulder shrugs/scapular retraction without resistance</li> </ul> </li> <li>Phase III: 7 - 9 Weeks         <ul> <li>Sling Immobilizer: May be discontinued</li> <li>Range of Motion: AROM in the pain free range - No PROM, AAROM (pulleys, supine wand, wall clin)</li></ul></li></ul>	Diagno	osis:	Date of Surg	gery:	
<ul> <li>Range of Motion: Supported pendulum exercises under guidance of PT</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, patient may ride stationary bike with operative arm in the sling</li> <li>Phase II: 4-6 Weeks</li> <li>Sling Immobilizer: Worn at all times - Sleep with pillow under elbow to support the operative arm Range of Motion: AAROM in the supine position with wand - Goal: Forward Flexion to 90 degrees</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, shoulder shrugs/scapular retraction without resistance</li> <li>Phase III: 7-9 Weeks</li> <li>Sling Immobilizer: May be discontinued</li> <li>Range of Motion: AROM in the pain free range - No PROM, AAROM (pulleys, supine wand, wall clin)         <ul> <li>Goals: Forward Flexion to 120 degrees and Abduction to 90 degrees, ER to tolerance, IR and Ension (wand behind the back)</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, submaximal isometrics</li> </ul> </li> <li>Phase IV: 10-12 Weeks         <ul> <li>Range of Motion: AROM and AAROM in the pain free range - No PROM, Goals: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone Sapular Retraction Exercises: (without weights), Wall Push-Ups (no elbow flexion &gt; 90 degrees)</li> </ul> </li> <li>Phase V: 3-6 Months         <ul> <li>Range of Motion: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May st t light weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups</li> <li>Return to athletics at 6 months post-op (no flies or pull downs), Regular Push-Ups</li> </ul> </li></ul>	Phase	I: 0 - 3 Weeks			
<ul> <li>Range of Motion: AAROM in the supine position with wand – Goal: Forward Flexion to 90 degrees</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, shoulder shrugs/scapular retraction without resistance</li> <li>Phase III: 7-9 Weeks</li> <li>Sling Immobilizer: May be discontinued</li> <li>Range of Motion: AROM in the pain free range – No PROM, AAROM (pulleys, supine wand, wall clin)         <ul> <li>Goals: Forward Flexion to 120 degrees and Abduction to 90 degrees, ER to tolerance, IR and Ension (wand behind the back)</li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, submaximal isometrics</li> </ul> </li> <li>Phase IV: 10-12 Weeks         <ul> <li>Range of Motion: AROM and AAROM in the pain free range – No PROM, Goals: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone Sapular Retraction Exercises (without weights), Wall Push-Ups (no elbow flexion &gt; 90 degrees)</li> </ul> </li> <li>Phase V: 3-6 Months         <ul> <li>Range of Motion: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May st t light weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups</li> <li>Return to athletics at 6 months postop</li> </ul> </li> <li>Comments:</li> </ul>	•	<b>Range of Motion:</b> Supported pendulum exercises under guidance of PT <b>Therapeutic Exercises:</b> Elbow and wrist active motion (with shoulder in neutral position at the side), hand exercises, patient may ride stationary bike with operative arm in the sling			
<ul> <li>Range of Motion: AROM in the pain free range - No PROM, AAROM (pulleys, supine wand, wall clin)         <ul> <li>Goals: Forward Flexion to 120 degrees and Abduction to 90 degrees, ER to tolerance, IR and Ension (wand behind the back)</li> </ul> </li> <li>Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side) hand exercises, submaximal isometrics</li> <li>Phase IV: 10-12 Weeks         <ul> <li>Range of Motion: AROM and AAROM in the pain free range - No PROM, Goals: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone Sapular Retraction Exercises (without weights), Wall Push-Ups (no elbow flexion &gt; 90 degrees)</li> </ul> </li> <li>Phase V: 3-6 Months         <ul> <li>Range of Motion: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May st t light weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups</li> <li>Return to athletics at 6 months postop</li> </ul> </li> <li>Comments:</li> </ul>	•	Range of Motion: AAROM in the supine position with wand – Goal: Forward Flexion to 90 degrees Therapeutic Exercises: Elbow and wrist active motion (with shoulder in neutral position at the side), hand exercises, shoulder shrugs/scapular retraction without resistance			
<ul> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone Sapular Retraction Exercises (without weights), Wall Push-Ups (no elbow flexion &gt; 90 degrees)</li> <li>Phase V: 3 - 6 Months</li> <li>Range of Motion: Full ROM</li> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May st t light weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups</li> <li>Return to athletics at 6 months postop</li> </ul> Comments:	•	Range of Motion: AROM in the pain free  Output  Output  Goals: Forward Flexion to 120 degension (wand behind the back)  Therapeutic Exercises: Elbow and wrist hand exercises, submaximal isometrics	rees and Abduction to 90 degr	ees, ER to tolerance, IR and Ex	
<ul> <li>Therapeutic Exercises: Light Theraband (ER, Abduction, Extension) with increasing resistance, May st t light weight training at 4.5 months post-op (no flies or pull downs), Regular Push-Ups</li> <li>Return to athletics at 6 months postop</li> </ul> Comments:	•	<b>Therapeutic Exercises:</b> Light Theraband (ER, Abduction, Extension), Biceps and Triceps PREs, Prone Sc apular Retraction Exercises (without weights), Wall Push-Ups (no elbow flexion > 90 degrees)			
	•	<b>Therapeutic Exercises:</b> Light Theraband t light weight training at 4.5 months post-order.	op (no flies or pull downs), Re	•	
Frequency: 2-3 times per week Duration: 6-8 weeks	Comm	ents:			
	Freque	ency: 2-3 times per week	Duration:	6-8 weeks	
Signature: Date:	Signature:		Date:		