

## Rehabilitation Protocol: Reverse Total Shoulder Replacement

Name: \_\_\_\_\_

Date: \_\_\_\_\_

Diagnosis: \_\_\_\_\_

Date of Surgery: \_\_\_\_\_

### Phase I (Weeks 0-2)

- Sling immobilization for first 2 week
  - Out of sling for home exercises (pendulums twice daily)
- Therapeutic Exercise
  - Grip Strengthening
  - Elbow/Wrist/Hand Exercises
  - Teach Home Exercises -- Pendulums
- Heat/Ice before and after PT sessions
- Limit External Rotation to Neutral (for Subscapularis Repair)

### Phase II (Weeks 3-12)

- Discontinue sling
- Range of Motion – PROM, AROM - increase as tolerated
  - Begin Active Internal Rotation and Backward Extension as tolerated
  - Goals: >90° Forward Flexion and 30° External Rotation
- Therapeutic Exercise
  - Begin light resisted exercises for Forward Flexion, External Rotation and Abduction – isometrics and bands – Concentric Motions Only
  - **No Resisted Internal Rotation, Backward Extension or Scapular Retraction**
  - Modalities per PT discretion

### Phase III (Months 3-12)

- Range of Motion – Progress to full AROM without discomfort – gentle passive stretching at end range
- Therapeutic Exercise
  - Begin resisted Internal Rotation and Backward Extension exercises
  - Advance strengthening as tolerated – Rotator Cuff, Deltoid and Scapular Stabilizers
  - Begin eccentric motions, plyometrics and closed chain exercises
- Modalities per PT discretion

Comments:

Frequency: 2-3 times per week

Duration: 6-8 weeks

Signature: \_\_\_\_\_

Date: \_\_\_\_\_