

Rehabilitation Protocol: High Tibial Osteotomy

name:	·	Date:	
Diagnosis:		Date of Surgery:	
Phase	e I: 1 - 5 Weeks		
• • • • Phase	Weightbearing ○ Toe-touch weightbearing with crutches and brace locked in extension (0-6 weeks) Hinged Knee Brace: Locked in extension for all activities (Including sleeping) – removed for PT Range of Motion: AROM/AAROM/PROM – Goal of 0-90 degrees under guidance of PT Therapeutic Exercises: Heel slides 0-90 degrees, quad sets, ankle pumps, non-weightbearing calf/hamstring stretches, straight leg raise with brace locked in full extension, resisted ankle dorsifl exion/ plantarflexion se II: 6-8 Weeks		
 Weightbearing Advance to 25% weightbearing with crutches each week and brace locked in extension Hinged Knee Brace: Unlocked for ambulation – May remove at night for sleeping Range of Motion: AROM/AAROM/PROM – Goal of 0-120 degrees under guidance of PT Therapeutic Exercises: Progress phase I exercises, straight leg raise with out of brace if capable of maintaining full extension, may add stationary bicycle No Closed Chain Activities Until 6 Weeks Post-op Phase III: 9-12 Weeks 			
•••Phase			
•	Weightbearing ○ Full weightbearing with a normal gait pattern Range of Motion: Full pain-free ROM Therapeutic Exercises: Progress closed chain activities, begin treadmill walking, swimming, and sp ort specific		
Commo	nents:		
Freque	nency: 2-3 times per week Dura	tion: 6-8 weeks	
Signatı	ture: Date	:	

