

## **Rehabilitation Protocol:**

Reverse Total Snoulder Replacement		
Name:		Date:
Diagnosis:		Date of Surgery:
Phase I (Weeks	s 0-2)	
Therape  O( O H  O H  Heat/Ice	Imobilization for first 2 week Out of sling for home exercises (pendulums eutic Exercise Grip Strengthening Elbow/Wrist/Hand Exercises Teach Home Exercises Pendulums e before and after PT sessions eternal Rotation to Neutral (for Subscapularis	
Phase II (Week	κs 3-12)	
Range of  Range of  Therape  Figure 1	inue sling of Motion – PROM, AROM - increase as tolera Begin Active Internal Rotation and Backwa Goals: >90° Forward Flexion and 30° Exter- eutic Exercise Begin light resisted exercises for Forward F and bands – Concentric Motions Only No Resisted Internal Rotation, Backward Modalities per PT discretion	rd Extension as tolerated nal Rotation Flexion, External Rotation and Abduction – isometrics
Phase III (Mont	ths 3-12)	
range Therape	of Motion – Progress to full AROM without deutic Exercise Begin resisted Internal Rotation and Backwadvance strengthening as tolerated – Rotatal Begin eccentric motions, plyometrics and coies per PT discretion	or Cuff, Deltoid and Scapular Stabilizers
Comments:		
Frequency: 2-3	3 times per week Dura	ation: 6-8 weeks





Signature:





Date:\_\_\_\_\_





